

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

wards at Woods Hole, Mass., U. S. National Museum No. 20247. Upon examining the fish in the container to which this number is given (labeled C. fallax), the writer found seven specimens, six of these were C. bartholomaei and one was C. latus. From the form of the fish, the lack of elevation of anterior rays of soft dorsal and anal; the greater number of soft rays (26 dorsal rays in drawing, 21 or 22 in latus; 22 anal rays in drawing, 16 to 18 in latus); the absence of a black blotch on opercle; the short maxillary, its tip not reaching anterior margin of pupil, and curved and straight portions of lateral line subequal, it is evident that the drawing was made from one of the specimens of C. bartholomaei.

Lewis Radcliffe, U. S. Bureau of Fisheries.

## DEKAY'S SNAKE IN THE LATE FALL

Previous to November 26, 1914, there were several drops of temperature to below the freezing point, 18 degrees above zero being reached on one occasion. On the above date, the weather was milder, the thermometer registering as high as 50 degrees. While the writer was walking near the corner of Kingsbridge Road and University Avenue, he found a young DeKay's snake (Storeria dekayi), about 8cm. in length. It was quite active and when released, quickly disappeared under the dried grass.

LEE S. CRANDALL, New York, N. Y.